## OSPHRANTER ROBUSTUS, Gould.

## Black Wallaroo.

Spec. Char.—Osph. artubus anticis magnis et prærobustis; vellere e fusco cinereo, infernè pallidiore; tarsis fuscis; digitis anticè nigris; antipedibus carpisque nigris; capite fuliginoso levitèr tincto; utrâque genâ lineâ albescente notatâ; gulâ, guttureque albidis; caudâ supernè fuscâ, subtùs pallidiore.

Descr.—Male. Fur harsh and somewhat shaggy; general colour slate-grey, obscurely washed with brownish, and tinted with vinous on the outer sides of the thighs; feet dark brown, gradually passing into black on the forepart; upper part of the arm brownish; hands and wrists black; inner surface of the ear white, the exterior brown; muzzle and a patch on the chin blackish; a line round the angle of the mouth and the lower lip white; throat and fore-part of the neck white, the hairs being grey at the base; under surface like the upper, but paler; tail blackish brown above, paler beneath.

Female. General colour silvery grey, obscurely tinted with purplish or vinous on the back; under surface nearly white; cheeks hoary, with a blackish patch on the chin; tail dirty white, slightly tinged with brown on the upper side; lcgs paler than the body; hands brown, becoming nearly black on the fingers; toes brownish black above.

|      |   |    |     |    |   | Male. |   |       |                   |  |   | Female. |                    |  |
|------|---|----|-----|----|---|-------|---|-------|-------------------|--|---|---------|--------------------|--|
|      |   |    |     |    |   |       |   | feet. | inches.           |  | 1 | feet.   | inches.            |  |
| Leng | th from the nose to the extremity of the tail |    |     |    |   |       |   | 7     | 0 .               |  |   | 5       | 10                 |  |
| ,,   | of tail                                       |    |     |    |   |       |   | 2     | $10\frac{1}{2}$ . |  |   | 2       | 6                  |  |
| ,,   | "tarsus and toes, including the nail          |    |     |    |   |       |   |       | 12 .              |  |   | 1       | $10^{\frac{1}{2}}$ |  |
| ,,   | ,, arm and hand, including the nails          |    |     |    |   |       |   |       | $13\frac{1}{2}$ . |  |   |         | $9\frac{1}{2}$     |  |
| ,,   | ,, face from the tip of the nose to the base  | of | the | ea | r |       |   |       | 8 .               |  |   |         | 7                  |  |
| ,,   | ,, ear  |    |     |    |   |       | • |       | $3\frac{1}{2}$ .  |  |   |         | 3                  |  |

Macropus (Petrogale) robustus, Gould in Proc. of Zool. Soc., Part VIII. p. 92. Black Wallaroo of the Colonists.

The Black Wallaroo inhabits the summits of sterile and rocky mountains, seldom descending to the coverts of their sides and never to their base; few, therefore, have had an opportunity of observing it in a state of nature; indeed there are thousands of persons in Australia who are not even aware of its existence. Although the south-eastern portion of the continent is, I believe, the only part of the country in which it has yet been observed, in all probability it has an extensive range northwards. It is tolerably abundant on the Liverpool range, and I ascertained that it inhabited many of those hills that branch off on either side of this great mountain-chain, both towards the interior as well as towards the coast. Its retreats are so well chosen among the crags and overhanging ledges of rocks, that it is nearly useless to attempt its capture with dogs. It is a formidable and even dangerous animal to approach, for if so closely pressed that it has no other chance of escape, it will rush at and force the invader over the edge of the rocks, as the Ibex is said to do under similar circumstances. Independently of its great muscular power, this animal is rendered still more formidable by the manner in which it makes use of its teeth, biting its antagonist with great severity.

The Black Wallaroo may be regarded as a gregarious animal, four, six and even more being frequently seen in company. On one of the mountains near Turi, to the eastward of the Liverpool Plains, it was very numerous; and from the nature of this and the other localities in which I observed it, it must possess the power of existing for long periods without water, that element being rarely to be met with in such situations.

The summits of the hills to which this species resorts soon become intersected by numerous roads and well-trodden tracks, caused by its repeatedly traversing from one part to the other; its food consists of grasses and the shoots and leaves of the low scrubby trees which clothe the hills it frequents.

Although much shorter in stature, and consequently less elegant in form, the fully adult male of this species equals in weight the largest specimens of *Macropus major*; and so remarkable is the difference in the colour and size of the sexes, that had I not seen them together in a state of nature, I should have considered them to be different species, the black and powerful male offering so great a contrast to the small and delicate female.